

A multiparameter assessment of the dose-dependent cytotoxicity of Camptothecin on HeLa cells

INTRODUCTION

High content cytotoxicity assays are in high demand for the cell-based analysis of anticancer drugs. In this study, we demonstrate an automated, multi-parameter assessment of dose-dependent Camptothecin cytotoxicity using the CELENA® X High Content Imaging System. The CELENA® X is a simple solution for the quantitative analysis of cytotoxicity because it can rapidly capture multi-channel fluorescence images from a multi-well plate and the integrated analysis software can accurately analyze multiple cellular parameters.

APPLICATION

Cell preparation

To study cytotoxicity *in vitro*, we treated HeLa cells with different concentrations of Camptothecin (CPT; Sigma-Aldrich, C9911). HeLa cells were counted with the LUNA-II™ Automated Cell Counter, seeded at a density of 1×10^4 HeLa cells/well on a 96-well plate, and cultured overnight. Cells were then treated with serial dilutions of Camptothecin (DMSO, 0.0625 μM , 0.125 μM , 0.25 μM , 0.5 μM , 1 μM , 2 μM , 4 μM , 8 μM , 16 μM , 32 μM , 64 μM) at four wells per condition with 95% humidity and 5% CO_2 at 37°C for 20 hours. To distinguish dead cells, cells were stained with propidium iodide (PI; Logos Biosystems, F23003). To assess total cell numbers, nuclei were counterstained with Hoechst 33342 (Life Technologies, H3570).

Imaging and analysis

Dead and total cells were visualized using the CELENA® X High Content Imaging System. Images were acquired using image-based autofocusing and a 4X LWD high NA objective in combination with filters for Hoechst 33342 (DAPI filter cube: Ex375/28, Em460/50) and propidium iodide (RFP filter cube: Ex530/40, Em605/55). One image field was acquired per well from 48 wells. Acquired images were analyzed using CELENA® X Cell Analyzer software.

For quantitative analysis of cytotoxicity, the integrated CELENA® X Cell Analyzer software was used to batch process and analyze images automatically. Two pipelines, *Cytotoxicity_DAPI_HeLa_4X* and *Cytotoxicity_RFP_HeLa_4X*, were established to determine the effects of Camptothecin (Table 1). Both pipelines, *Cytotoxicity_DAPI_HeLa_4X* and *Cytotoxicity_RFP_HeLa_4X*, employed the *IdentifyPrimaryObjects* module to identify individual cells based on Hoechst 33342 or propidium iodide staining, respectively. Both used the *MeasureImageAreaOccupied* module measured the surface area occupied by the fluorescence-stained cells. In the *Cytotoxicity_RFP_HeLa_4X* pipeline, propidium iodide-stained cells were further analyzed with the *MeasureObjectSizeShape* module to quantify their size and shape, followed by the *MeasureObjectIntensity* module to measure propidium iodide signal intensity.

Table 1. Pipelines used to assess transfection efficiency with CELENA® X Cell Analyzer

Pipeline	Order	Module	Use
Cytotoxicity_DAPI_HeLa_4X	1	IdentifyPrimaryObjects	To find Hoechst-stained nuclei
	2	MeasureImageAreaOccupied	To quantify the surface area occupied by the Hoechst-stained nuclei
	3	GrayToColor	To produce color images from grayscale images
	4	OverlayOutlines	To place outlines around identified nuclei
Cytotoxicity_RFP_HeLa_4X	1	IdentifyPrimaryObjects	To find PI-stained nuclei
	2	MeasureImageAreaOccupied	To quantify the surface area occupied by the PI-stained nuclei
	3	MeasureObjectSizeShape	To measure the size and shape of the PI-stained nuclei
	4	MeasureObjectIntensity	To measure RFP intensity
	5	GrayToColor	To produce color images from grayscale images
	6	OverlayOutlines	To place outlines around identified nuclei

To visualize the data, both pipelines used the *GraytoColor* module to apply pseudocolor to the grayscale images of each channel and then the *OverlayOutlines* module to outline the identified cells. GraphPad Prism® software (GraphPad Software, Inc.) was used to plot sigmoidal dose-response curves and calculate IC₅₀ values.




Figure 1. Hoechst-stained HeLa cells after 20 hours of Camptothecin treatment. (A) Image analysis of Camptothecin-induced cytotoxicity with green borders outlining Hoechst-stained nuclei. Dose-response curves show the cytotoxic effects of Camptothecin on Hoechst-determined (B) cell count and (C) occupied surface area.

Figure 1 showed that higher doses of Camptothecin exposure decreased the number of Hoechst-stained nuclei and the area that they occupied. The IC₅₀ of Camptothecin in relation to these two parameters was 5.2180 μ M on average. Figure 2 shows that higher Camptothecin concentrations lead to an increase in clumpy cellular aggregates (Fig. 2A, B), propidium iodide signal size (Fig. 2C), and signal intensity (Fig. 2D). These data demonstrate that cell death leads to the formation of clumpy aggregates caused by the presence of cell debris and free DNA from dead cells. The IC₅₀ of Camptothecin upon assessing these parameters was 4.8591 μ M on average. Combined, these results indicate a Camptothecin dose-dependent increase of cell death.




Figure 2. Propidium iodide-stained HeLa cells after 20 hours of Camptothecin treatment. (A) Image analysis of Camptothecin-induced cytotoxicity with green borders outlining propidium iodide-stained nuclei. Dose-response curves show the cytotoxic effects of Camptothecin on (B) occupied surface area, (C) size, and (D) integrated intensity of the propidium iodide signal.

CONCLUSION

In this study, we used the CELENA[®] X to evaluate the dose-dependent cytotoxicity of Camptothecin in a simple, effective, and highly reproducible way. The CELENA[®] X allowed us perform a multi-parameter assessment on the effects of Camptothecin on cell viability. Using the CELENA[®] X High Content Imaging System and CELENA[®] X Cell Analyzer software, the same experimental conditions as well as the identical analysis pipeline can be reused to verify results.

REFERENCES

- Peter S, Yu H, Ivanyi-Nagy R, Dröge P. 2016. Cell-based high-throughput compound screening reveals functional interaction between oncofetal HMGA2 and topoisomerase I. *Nucleic Acids Research*, 44(22): e162.
- Zhang L, Ma D, Zhang Y, He W, Yang J, Li C, Jiang H. 2013. Characterization of DNA topoisomerase-1 in *Spodoptera exigua* for toxicity evaluation of camptothecin and hydroxy-camptothecin. *PLoS One*, 8(2): e56458.